

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/302959005>

Hepatoprotective properties of Dandelion: Recent update

Article in *Journal of Applied Pharmaceutical Science* · May 2016

DOI: 10.7324/JAPS.2016.60429

CITATIONS

18

READS

6,981

1 author:



[Ezhilarasan Devaraj](#)

Saveetha University

163 PUBLICATIONS 2,319 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Collaborative Book Project on Phytochemistry, Molecular Biology & Biochemistry, Nutrition & Toxicology [View project](#)



Wnt signaling modulators in ALD and NAFLD [View project](#)

Hepatoprotective properties of Dandelion: recent update

Ezhilarasan Devaraj

Department of Pharmacology, Saveetha Dental College and Hospitals Saveetha University, Chennai - 600 077, Tamil Nadu, India.

ARTICLE INFO

Article history:

Received on: 24/10/2015

Revised on: 20/12/2015

Accepted on: 10/02/2016

Available online: 30/04/2016

Key words:

Taraxacum officinale,
oxidative stress, antioxidant,
hepatoprotective

ABSTRACT

Taraxacum officinalis (Dandelion) have been used as folk medicines in China, India, and Russia for the treatment of chronic liver diseases. This review focuses exclusively on published literature pertaining to the potential use of *Taraxacum officinalis* or Dandelion for the treatment of various chronic liver diseases. Dandelion has been tested against various drugs and chemical induced chronic liver diseases in experimental animals and came out with promising results. *In vitro* studies also confirm the hepatoprotective, antioxidant and anti-inflammatory properties of dandelion. The extensive literature survey reveal the fact that *Taraxacum officinalis* or dandelion to be safe and the available evidence on the mechanisms of action appears promising, there are currently insufficient data from well-conducted clinical trials to recommend it use in patients with alcoholic liver chronic liver disease.

INTRODUCTION

The liver is one of the vital organs of human body, being involved in myriad vital functions and regulation of physiological processes. Due to its unique anatomic location and function, liver is vulnerable to many forms of injury. Fortunately, it also has remarkable capacity to repair itself following many types of injuries that it encounters in compensation for the important, yet hazardous functions it undertakes. However, liver disorders following chronic liver injury lead to various pathological changes such as fatty liver, increase in ROS or oxidative stress, necrosis of liver cells, hepatitis, steatosis, cholestasis, vascular lesions, and granuloma and veno-occlusive diseases, increase in the level of inflammatory markers, fibrosis, cirrhosis, hepatocellular carcinoma which further produce portal hypertension and organ failure (Singh *et al.*, 2011).

The conventional treatments of chronic liver injuries such as fibrosis, cirrhosis, steatosis and chronic hepatitis are frequently inadequate due to side effects caused by various drugs and chemicals. To overcome this problem, current research focused on drugs derived from medicinal plants, due to their richness in flavonoid and polyphenolic compounds have been extensively studied against various drug and chemical induced

hepatotoxicity *in vivo* and *in vitro* and it is need of the hour (Pereira *et al.*, 2015). There are several plant derived agents have shown promising hepatoprotective properties *i.e.*, dandelion, silymarin, silibinin, curcumin, berberin, resveratrol (Ezhilarasan *et al.*, 2014; Ezhilarasan *et al.*, 2012). Among these dandelion is one of the promising agent obtained from the medicinal plant *Taraxacum officinalis*. Hence, the present review aimed to gain insight into the recent update on the hepatoprotective properties of *Taraxacum officinalis*.

Description

The genus *Taraxacum* Wiggers, family Asteraceae, subfamily Cichorioideae, tribe Lactuceae, commonly known as dandelion, includes approximately 30–57 varieties with many microspecies, divided into nine sections (Vašut *et al.*, 2015). *Dandelion* (*Taraxacum officinalis*), a traditional herbal medicine is used for treatment of jaundice and disorders of the liver, gallbladder and other various hepatic ailments (You *et al.*, 2010; Ahmed *et al.*, 2013,). The folk medicines of China, India, and Russia have recognized dandelion's effect as a liver tonic. Traditional Chinese medicine combines dandelion with other herbs to treat hepatitis (Modaresi, 2012). Conventionally, root and herb from *Taraxacum officinale* (TO) have been reported to use for the treatment of various ailments, including liver and gallbladder disorders (Gulfranz *et al.*, 2014).

* Corresponding Author
Email: ezhild@gmail.com

Phytochemical constituents

The phytochemical investigation showed that TO has an abundance of terpenoid and sterol (principally taraxacin and taraxacerin), equally distributed in the roots, leaves, and flowers. Other terpene/sterol compounds include beta-amyrin, taraxasterol, and taraxerol, as well as free sterols (sitosterin, stigmasterin, and phytosterin) structurally related to bile (Koo *et al.*, 2004; Schütz *et al.*, 2006). Presence of these biologically active principles in the TO extract have been attributed for its beneficial effects. The terms *Taraxacum officinale* and dandelion are generally interchangeably used in the literature.

SPECTRUM OF HEPATOPROTECTIVE EFFECTS

Antioxidant effect

Oxidative stress is a common feature observed in a wide spectrum of chronic liver diseases including viral hepatitis, alcoholic, and nonalcoholic steatohepatitis (Jiang and Török, 2015). Oxidative stress leads to deleterious processes in the liver and produces liver diseases. Therefore, restoring antioxidants is essential to maintain homeostasis. One method of restoring antioxidants is suggested to consume natural compounds with antioxidant capacity (Casas-Grajales and Muriel, 2015). Dandelion, a natural antioxidant compound has been empirically used due to its health-promoting properties as an anti-carcinogenic, anti-inflammatory and anti-oxidant (You *et al.*, 2010). Ethanolic TO leaves and root extract significantly attenuated marker enzymes of liver toxicity (AST and ALT), lipid peroxidation and oxidative stress induced by acetaminophen in mice. The TO extract shown to possess the free radical quenching activities. This protective effects of TO have been suggested to the presence of phenolic compounds in the extract (Colle *et al.*, 2012). In a recent study, ethanolic and n-hexane TO leaves extract significantly decreased the liver marker enzymes, superoxide dismutase (SOD), catalase, lipid peroxidation and glutathione

peroxidase (GPx) in rats intoxicated with CCl₄. The efficacy of TO ethanolic leaves extract found to be more effective as it compared with n-hexane extract and silymarin against CCl₄ induced hepatotoxicity and oxidative stress in rats (Gulfraz *et al.*, 2014).

In CCl₄ induced oxidative stress model TO extract reversed the glutathione (GSH) depletion, up-regulation of Nuclear factor-kB (NF-kB) and increased expression of regulatory inflammatory mediators, such as inducible nitric oxide synthase (iNOS), cyclooxygenase (COX)-2, Tumor necrosis factor- α (TNF- α) and interleukin (IL)-1 α (Park *et al.*, 2010). These results suggest that TO have a hepatoprotective effect by modulating inflammatory responses and ameliorating oxidative stress. Alcohol is a leading cause of liver disease and is associated with significant morbidity and mortality. Alcohol induced liver diseases represents a spectrum of liver pathology ranging from fatty change to fibrosis to cirrhosis (Dugum and McCullough, 2015). *In vitro* the protective effects of TO root against alcoholic liver damage were investigated in HepG2/2E1 cells. In this study, ROS was generated by the administration of ethanol *in vitro* this was consequent to decrease in cell viability by less than 40%. However, cells were simultaneously treated with ethanol and TO hot water root extract did not induce cytotoxicity as compare to ethanol alone treated HepG2/2E1 cells (You *et al.*, 2010).

Hu and Kitts (2004) reported that dandelion flower extracts has also the capability to scavenge ROS and prevent DNA from ROS-induced damage *in vitro*. Suppression of oxidative stress by dandelion has been attributed to presence of luteolin and luteolin-7-Oglucoside. Generally, *Taraxacum* species are found effective against oxidative stress. Recently, it has been reported that *T. marginellum* was the most efficient extract reducing intracellular ROS levels although *in vitro* assays, *T. obovatum* was observed as the best free radical scavenger (Mingarro *et al.*, 2015). Undoubtedly the above studies suggest the promising efficacy of TO against various drugs induced oxidative stress.




Fig. 1: Hepatoprotective effects of *Taraxacum officinale*.

Antifibrotic effects

Hepatic fibrosis is common sequel following chronic liver injury and reversal of fibrosis prior it attains the cirrhotic stage would be a clear therapeutic strategy (Lee *et al.*, 2015). TO extract has been evaluated against the drug and chemical induced hepatic fibrosis in experimental animals and came out with promising results. For instance, it has been reported that dandelion root water-ethanolic extract (DWE) ameliorated the CCl₄ induced hepatic fibrosis in mice. Administration of 600 mg/kg of DWE for 10 days in mice shows significant replenishment of liver aspartate and alanine transaminases (AST and ALT) marker enzymes of hepatotoxicity, superoxide dismutase, hydroxyproline and α -smooth muscle actin (α -SMA) protein expression in CCl₄ induced hepatic fibrotic mice (Domitrović *et al.*, 2010). Further, this study suggests that administration TO promote the complete regression of fibrosis and the enchantment of hepatic regenerative capabilities.

Antisteatotic effect

Steatosis or alcoholic fatty liver disease has a widespread incidence and is the first step in the progression to more severe stages of alcoholic liver disease, with concomitant increases in morbidity and mortality rates (Lívero and Acco, 2015). CCl₄ induced lipid changes in liver is one of the classical model for the evaluation of lipid lowering activity of drug and herbal extract. Ethanol and n-hexane leaves extract of TO significantly lower the lipid profile in CCl₄ administered rats. Interestingly, TO significantly reduces the increase in triglyceride (TG), cholesterol (CHO), high and low density lipoproteins (HDL and LDL) (Gulfranz *et al.*, 2014). It is reported that dandelion leaves extract has the ability to reduce the murine model of methionine- and choline-deficient diet-induced nonalcoholic steatohepatitis (NASH). In light of these studies it has been suggested that TO could also be a promising agent and has to be tested against hepatic steatosis condition (Davaatseren *et al.*, 2013). In a recent study TO leaves extract found effective against CCl₄ induced hepatic steatosis in rats. It was shown that steatosis grade was significantly reduced upon administration of TO leave extract (Al-Malki *et al.*, 2013)

Anticancer activity

Hepatocellular carcinoma (HCC) is one of the most common malignancies, which accounts for 90% of primary liver cancer. HCC usually presents with poor outcomes due to the high rates of tumor recurrence and widespread metastasis (Mao and Wang, 2015). Recent results pointed out that natural products, in particular those present in *Taraxacum* root extract, have great potential as non-toxic and effective alternatives to conventional modes of chemotherapy available today (Ovadje *et al.*, 2012). Dandelion has been said to induce cytotoxicity in Hep G2 cells and decreases its viability below 40% (You *et al.*, 2010). TO extract significantly induced the secretion of TNF- α and IL-1 α and apoptosis of Hep G2 cells (Koo *et al.*, 2004). These strategies are clearly implicated in anticancer efficacy of dandelion. In a very

recent study it was found that relevant cytotoxic effect in *T. lacistrum* extract over HeLa and HepG2 cell lines (Mingarro *et al.*, 2015). However, studies regarding the usefulness of TO against liver carcinoma experimental models are scanty or not available in literature. Hence, further studies are warranted in animal models to prove the promising anti-cancer efficacies of TO observed *in vitro* cancer cell line models *i.e.*, Hep G2 and HeLa.

CONCLUSION

Overwhelming evidence clearly suggest that dandelion is widely used in traditional and natural medicine systems worldwide. Thus far it has received little research attention towards its beneficial effects against various hepatic ailments. In light of the studies reviewed here it has been evident that TO has the ability to intervene various pathophysiological functions related to liver. The extensive literature survey reveal the fact that *Taraxacum officinale* or dandelion appear to be safe and the available evidence on the mechanisms of action appears promising, there are currently insufficient data from well-conducted clinical trials to recommend their use in patients with chronic liver diseases.

REFERENCES

- Ahmed D, Gulfranz M, Ahmad MS, Tahir RM, Anwar P. Cytoprotective potential of methanolic leaves extract of *Taraxacum officinale* on CCl₄ induced Rats. *Pensee J* 2013; 75:220-227.
- Al-Malki AL, Abo-Golayel MK, Abo-Elnaga G, Al-Beshri H. Hepatoprotective effect of dandelion (*Taraxacum officinale*) against induced chronic liver cirrhosis. *J Med Plants Res* 2013; 7:1494-1505
- Casas-Grajales S, Muriel P. Antioxidants in liver health. *World J Gastrointest Pharmacol Ther* 2015; 6:59-72.
- Colle D, Arantes LP, Gubert P, da Luz SC, Athayde ML, Teixeira Rocha JB, Soares FA. Antioxidant properties of *Taraxacum officinale* leaf extract are involved in the protective effect against hepatotoxicity induced by acetaminophen in mice. *J Med Food* 2012; 15(6):549-556.
- Davaatseren M, Hur HJ, Yang HJ, Hwang JT, Park JH, Kim HJ, Kim MJ, Kwon DY, Sung M. *Taraxacum officinale* (dandelion) leaf extract alleviates high-fat diet-induced nonalcoholic fatty liver. *Food Chem Toxicol* 2013; 58:30-36.
- Domitrović R, Jakovac H, Romić Z, Rahelić D, Tadić Z. Antifibrotic activity of *Taraxacum officinale* root in carbon tetrachloride-induced liver damage in mice. *J Ethnopharmacol* 2010; 130:569-577.
- Dugum M, McCullough A. Diagnosis and Management of Alcoholic Liver Disease. *J Clin Transl Hepatol* 2015; 3:109-116.
- Ezhilarasan D, Karthikeyan S, Vivekanandan P. Ameliorative effect of silibinin against *N*-nitrosodimethylamine-induced hepatic fibrosis in rats. *Environ Toxicol Pharmacol* 2012; 34:1004-1013.
- Ezhilarasan D, Sokal E, Karthikeyan S, Najimi M. Plant derived antioxidants and antifibrotic drugs: Past, Present and Future. *J Coast Life Med* 2014; 2:738-745.
- Gulfranz M, Ahmad D, Ahmad MS, Qureshi R, Mahmood RT, Jabeen N, Abbasi KS. Effect of leaf extracts of *Taraxacum officinale* on CCl₄ induced hepatotoxicity in rats, *in vivo* study. *Pak J Pharm Sci* 2014; 27:825-829.
- Hu C, Kitts DD. Luteolin and luteolin-7-O-glucoside from dandelion flower suppress iNOS and COX-2 in RAW264.7 cells. *Mol Cell Biochem* 2004; 265:107-113.
- Jiang JX, Török NJ. NADPH Oxidases in Chronic Liver Diseases. *Adv Hepatol* 2014 Nov30; 2014.

Koo HN, Hong SH, Song BK, Kim CH, Yoo YH, Kim HM. *Taraxacum officinale* induces cytotoxicity through TNF-alpha and IL-1alpha secretion in Hep G2 cells. *Life Sci* 2004; 74:1149-1157.

Lee YA, Wallace MC, Friedman SL. Pathobiology of liver fibrosis: a translational success story. *Gut* 2015; 64:830-841.

Lívero FA, Acco A. Molecular basis of alcoholic fatty liver disease: From incidence to treatment. *Hepatol Res* 2015 Sep 29. doi: 10.1111/hepr.12594.

Mao B, Wang G. MicroRNAs involved with hepatocellular carcinoma (Review). *Oncol Rep*. 2015 Sep 15. doi: 10.3892/or.2015.4275.

Mingarro DM, Plaza A, Galán A, Vicente JA, Martínez MP, Acero N. The effect of five *Taraxacum* species on *in vitro* and *in vivo* antioxidant and antiproliferative activity. *Food Funct* 2015; 6:2787-2793.

Modaresi M, Resalatpour N. The Effect of *Taraxacum officinale* Hydroalcoholic Extract on Blood Cells in Mice. *Adv Hematol* 2012; 2012:653412.

Ovadge P, Chochkeh M, Akbari-Asl P, Hamm C, Pandey S. Selective induction of apoptosis and autophagy through treatment with dandelion root extract in human pancreatic cancer cells. *Pancreas* 2012; 41:1039-1047.

Park CM, Youn HJ, Chang HK, Song YS. TOP1 and 2, polysaccharides from *Taraxacum officinale*, attenuate CCl(4)-induced hepatic damage through the modulation of NF-kappaB and its regulatory mediators. *Food Chem Toxicol* 2010; 48:1255-1261.

Pereira C, Barros L, Ferreira IC. Extraction, identification, fractionation and isolation of phenolic compounds in plants with hepatoprotective effects. *J Sci Food Agric* 2015. doi: 10.1002/jsfa.7446.

Schütz K, Muks E, Carle R, Schieber A. Separation and quantification of inulin in selected artichoke (*Cynara scolymus* L.) cultivars and dandelion (*Taraxacum officinale* WEB. ex WIGG.) roots by high-performance anion exchange chromatography with pulsed amperometric detection. *Biomed Chromatogr* 2006; 20:1295-1303.

Singh A, Bhat TK, Om PS. Clinical Biochemistry of Hepatotoxicity. *J Clin Toxicol* S4:001.

Vašut R., Majeský L. *Taraxacum pudicum*, a new apomictic microspecies of T. section *Erythrosperma* (Asteraceae) from Central Europe. *Phytotaxa* 2015; 227:243–252.

You Y, Yoo S, Yoon HG, Park J, Lee YH, Kim S, Oh KT, Lee J, Cho HY, Jun W. In vitro and in vivo hepatoprotective effects of the aqueous extract from *Taraxacum officinale* (dandelion) root against alcohol-induced oxidative stress. *Food Chem Toxicol* 2010; 48:1632-1637.

How to cite this article:

Ezhilarasan Devaraj. Hepatoprotective properties of Dandelion: recent update. *J App Pharm Sci*, 2016; 6 (04): 202-205.